

Balan	Kerb	Prefar	
●	○	●	Summer Grasses
●	○	●	Pigweed
●	●	●	Purslane
●	●	○	Goundcherry
●	●	●	Goosefoot
●	●	●	Lambquarters
○	●	○	London rocket
○	●	○	Shepards purse
○	○	○	Sowthistle
●	●	●	Knotweed
●	●	○	Cereals
●	●	○	Wild oat
●	●	●	Canarygrass
●	●	●	Bluegrass

● = Excellent

● = Partial

○ = None



RESTRICTED USE PESTICIDE



KERB 50-W

SELECTIVE HERBICIDE IN WATER-SOLUBLE POUCHES

ACTIVE INGREDIENT
Pronamide

3,5-dichloro-N-(1,1-dimethyl-2-propynyl)-benzamide 51.0%

INERT INGREDIENTS 49.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

NOTES: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and disposal instructions on back panel and attached Use Directions. If the Conditions of Sale and Warranty are not acceptable, return the product within 30 days of purchase to the place of purchase.



INDEPENDENCE MALL WEST
PHILADELPHIA, PA. 19106-2389
PHONE (215) 592-3000


EPA Reg. No. 707-159
EPA Est. No. 65387-AR-001 (H)
Superscript used is first letter of Lot Number.

NET CONTENTS: 3 LBS (1.36 KG)
3 x 1 LB WATER-SOLUBLE POUCHES

MADE IN ITALY



Kerb (new)
1.5
heavy sprinklers

A photograph of a field with a new 1.5m furrow. The furrow is a long, straight, narrow trench in the soil, filled with dark, moist earth. The soil on either side of the furrow is light brown and appears dry, with some small green plants growing in the furrow. A yellow marker is visible in the background, and a white label with text is placed in the foreground.

Kerb (new)
1.5
furrow










Furrow



Sprinkler

Delayed Application

- Apply after sprinklers have started and before weeds are well established to avoid leaching the herbicide

Delayed Application


- After sprinklers have started, there are only two options for application:
 - Aerial application
 - Chemigation through sprinklers

Hours to Germination

- Summer annual weeds: 12-72
- Winter annual weeds: 72-168



**October 27- November 3
1999**





Untreated



Applied before
sprinklers started








Applied 3 days
after
sprinklers started








**November 2-8
2000**



**September 1-7
1999**





PRE-
SPRINKLERS




3 DAYS





14 DAYS

**September 12-18
2000**




**January 18-24
2001**



A photograph of a field of green plants, likely a crop field, with a white label in the foreground that reads "Untreated". The plants are arranged in rows, and the soil is visible between them. A small pink flag is visible in the background. The label is placed on the ground, and the plants are growing in the background.

Untreated



Pre-sprinklers

6 Days





Problems With This Strategy

- Application is less precise in rate and placement
- Overlaps and skips
- Increased weed control=increased crop injury







THE END